

AN INEQUALITY FOR BASIC CONFLUENT HYPERGEOMETRIC SERIES

RUIMING ZHANG

ABSTRACT. In this very short note we will derive an inequality for a class of entire functions including all the confluent basic hypergeometric series and an inequality for a class of meromorphic functions including theta functions.

1. INTRODUCTION

In the studies of Plancherel-Rotach type asymptotics of q -orthogonal polynomials such as Ismail-Masson $\{h_n(x|q)\}_{n=0}^{\infty}$, Stieltjes-Wigert $\{S_n(x; q)\}_{n=0}^{\infty}$ and q -Laguerre $\{L_n^{(\alpha)}(x; q)\}_{n=0}^{\infty}$ [4, 9, 10], the Ramanujan function $A_q(z)$, which is also called q -Airy function in the literature, plays a very important role. It is called q -Airy function because it appears repeatedly in the Plancherel-Rotach type asymptotics of q -orthogonal polynomials, just like classical Airy function in the classical Plancherel-Rotach asymptotics of classical orthogonal polynomials [5, 12]. In [14], we have investigated certain properties of the Ramanujan function $A_q(z)$. It turns out that, the polynomials like Ismail-Masson $\{h_n(x|q)\}_{n=0}^{\infty}$, Stieltjes-Wigert $\{S_n(x; q)\}_{n=0}^{\infty}$, q -Laguerre $\{L_n^{(\alpha)}(x; q)\}_{n=0}^{\infty}$ and the Ramanujan function $A_q(z)$, they are all in a particular class of entire functions including all the confluent basic hypergeometric series. In this short note will show that this class of entire functions satisfy an inequality similar to the inequality for the Ramanujan function $A_q(z)$ in [14]. Finally, we will state an inequality for a class of meromorphic functions including $\Theta(z|q)$ defined as in [9, 10]. $\Theta(z|q)$ also appears in the main terms of the Plancherel-Rotach type asymptotics when a scaling parameter is irrational.

As in papers [9, 10, 14], we will follow the usual notations from q -series [1, 2, 5]

$$(1) \quad (a; q)_0 := 1 \quad (a; q)_n := \prod_{k=0}^n (1 - aq^k), \quad \left[\begin{matrix} n \\ k \end{matrix} \right]_q := \frac{(q; q)_n}{(q; q)_k (q; q)_{n-k}}.$$

Moreover, we shall use the multishifted factorials [1, 2, 5]

$$(2) \quad (a_1, a_2, \dots, a_k; q)_n := \prod_{j=1}^k (a_j; q)_n.$$

Throughout this paper, we shall always assume that $0 < q < 1$, hence $n = \infty$ is allowed in the above definitions.

1991 *Mathematics Subject Classification*. Primary 30E15. Secondary 33D45.

Key words and phrases. Basic confluent hypergeometric series, q -Airy function, Ramanujan's entire function, inequalities, entire functions, meromorphic functions, theta functions.

A confluent basic hypergeometric series is formally defined as [1, 2, 5]

$$(3) \quad {}_r\phi_s \left(\begin{matrix} a_1, \dots, a_r \\ b_1, \dots, b_s \end{matrix} \middle| q, z \right) = \sum_{k=0}^{\infty} \frac{(a_1, \dots, a_r; q)_k}{(b_1, \dots, b_s; q)_k} z^k \left(-q^{(k-1)/2} \right)^{k(s+1-r)}$$

with $s+1-r > 0$.

We will use a simple summation formula

$$(4) \quad \frac{1}{(z; q)_{\infty}} = \sum_{k=0}^{\infty} \frac{z^k}{(q; q)_k},$$

which is a special case of the q -binomial theorem [1, 2, 5],

$$(5) \quad \frac{(az; q)_{\infty}}{(z; q)_{\infty}} = \sum_{k=0}^{\infty} \frac{(a; q)_k}{(q; q)_k} z^k.$$

The Ramanujan function $A_q(z)$ is defined as [5]

$$(6) \quad A_q(z) := \sum_{k=0}^{\infty} \frac{q^{k^2}}{(q; q)_k} (-z)^k.$$

In [14] we proved that

$$(7) \quad |A_q(z)| \leq \frac{\left(\frac{|z|}{\sqrt{q}} \right)^{1/2} \exp \left\{ -\frac{\log^2 |z|}{4 \log q} \right\}}{(q; q)_{\infty}}$$

for any nonzero complex number z and

$$(8) \quad |A_q(z)| \leq e^{q|z|/(1-q)}$$

for any complex number z .

The theta function $\Theta(z|q)$ is defined as

$$(9) \quad \Theta(z|q) = \sum_{k=-\infty}^{\infty} q^{k^2} z^k,$$

for $0 < q < 1$ and all nonzero complex number z .

2. THE INEQUALITY

Let

$$(10) \quad f(z) = \sum_{k=0}^{\infty} \frac{(a_1, \dots, a_r; q)_k q^{lk^2}}{(b_1, \dots, b_s, q; q)_k} z^k,$$

with the parameters satisfying

$$(11) \quad 0 \leq b_1, \dots, b_s < 1 \quad l > 0.$$

It is clear that the function

$$(12) \quad {}_r\phi_s \left(\begin{matrix} a_1, \dots, a_r \\ b_1, \dots, b_s \end{matrix} \middle| q, -zq^{(s+1-r)/2} \right)$$

is of the form (10) with

$$(13) \quad l = \frac{s+1-r}{2}.$$

Assume

$$(14) \quad c := c(a_1, \dots, a_r; b_1, \dots, b_s; q) = \frac{(-|a_1|, \dots, -|a_r|; q)_\infty}{(b_1, \dots, b_s; q)_\infty},$$

then,

$$(15) \quad |f(z)| \leq c \sum_{k=0}^{\infty} \frac{q^{lk^2} |z|^k}{(q; q)_k}.$$

We rewrite (15) as

$$(16) \quad c \sum_{k=0}^{\infty} \frac{q^{kl} (q^{l(k-1)} |z|)^k}{(q; q)_k}.$$

For $k = 0, 1, \dots$, the terms

$$(17) \quad \left(q^{l(k-1)} |z| \right)^k$$

are bounded by

$$(18) \quad \sqrt[4]{|z|^2 q^{-l}} \exp \left(-\frac{\log^2 |z|}{4l \log q} \right),$$

thus,

$$(19) \quad |f(z)| \leq c \sqrt[4]{|z|^2 q^{-l}} \exp \left(-\frac{\log^2 |z|}{4l \log q} \right) \sum_{k=0}^{\infty} \frac{q^{kl}}{(q; q)_k},$$

or

$$(20) \quad |f(z)| \leq \frac{c}{(q^l; q)_\infty} \sqrt[4]{|z|^2 q^{-l}} \exp \left(-\frac{\log^2 |z|}{4l \log q} \right).$$

Theorem 2.1. Assume that the entire function $f(z)$ is defined as in (10) and (11), then,

$$(21) \quad |f(z)| \leq \frac{(-|a_1|, \dots, -|a_r|; q)_\infty}{(b_1, \dots, b_s, q^l; q)_\infty} \left\{ \sqrt[4]{|z|^2 q^{-l}} \exp \left(-\frac{\log^2 |z|}{4l \log q} \right) \right\}$$

for any nonzero complex number z . In particular, for

$$(22) \quad 0 \leq b_1, \dots, b_s < 1, \quad 0 < s + 1 - r,$$

we have

$$(23) \quad \left| {}_r\phi_s \left(\begin{matrix} a_1, \dots, a_r \\ b_1, \dots, b_s \end{matrix} \middle| q, z \right) \right| \leq \frac{(-|a_1|, \dots, -|a_r|; q)_\infty}{(b_1, \dots, b_s, q^{(s+1-r)/2}; q)_\infty} \\ \times \left\{ \sqrt[4]{|z|^2 q^{3(r-s-1)/2}} \exp \left(\frac{\log^2 [|z| q^{(r-s-1)/2}]}{2(r-s-1) \log q} \right) \right\}$$

for any nonzero complex number.

The same proof could be employed to show the following results.

Theorem 2.2. *Let $f(z)$ be a meromorphic function with the following Laurent expansion*

$$(24) \quad f(z) = \sum_{k=-\infty}^{\infty} a_k (z-a)^k$$

on the whole complex plane except at $z = a$. If there exist two positive numbers $\alpha > 0$ and $0 < q < 1$ such that

$$(25) \quad c = \sum_{k=-\infty}^{\infty} |a_k| q^{-|k|^{\alpha+1}} < \infty,$$

then

$$(26) \quad |f(z)| \leq c \exp \{ \beta [|\log |z-a||]^{\gamma} \},$$

for $z \neq a$, where

$$(27) \quad \beta = \frac{\alpha}{(\alpha+1)^{1+1/\alpha} \log^{1/\alpha} q^{-1}},$$

and

$$(28) \quad \gamma = \frac{\alpha+1}{\alpha}.$$

In the particular, the theta function $\Theta(z|q)$ has an upper bound

$$(29) \quad |\Theta(z|q)| \leq c \exp \left(\frac{|\log |z||^2}{\log q^{-1}} \right)^{1/\alpha}$$

for any $0 < \alpha < 1$, where

$$(30) \quad c = \sum_{k=-\infty}^{\infty} q^{k^2 - |k|^{1+\alpha}}.$$

Proof. Clearly, we have

$$(31) \quad |f(z)| \leq \sum_{k=-\infty}^{\infty} |a_k| q^{-|k|^{\alpha+1}} \left(q^{|k|^{\alpha+1}} |z-a|^k \right).$$

For $k \in \mathbb{Z}$ and $z \neq a$, then, all the terms

$$(32) \quad q^{|k|^{\alpha+1}} |z-a|^k$$

are bounded by

$$(33) \quad \exp \frac{\alpha [|\log |z-a||]^{1+1/\alpha}}{(\alpha+1)^{1+1/\alpha} \log^{1/\alpha} q^{-1}},$$

and the inequality (26) follows. $\square$

REFERENCES

- [1] G. E. Andrews, R. A. Askey, and R. Roy, Special Functions, Cambridge University Press, Cambridge, 1999.
- [2] G. Gasper and M. Rahman, Basic Hypergeometric Series, second edition Cambridge University Press, Cambridge, 2004.
- [3] W. K. Hayman, On the zeros of a q -Bessel function, Contemporary Mathematics, volume 382, American Mathematical Society, Providence, 2005, 205–216.
- [4] M. E. H. Ismail, Asymptotics of q -orthogonal polynomials and a q -Airy function, Internat. Math. Res. Notices 2005 No 18 (2005), 1063–1088.
- [5] M. E. H. Ismail, Classical and Quantum Orthogonal Polynomials in one Variable, Cambridge University Press, Cambridge, 2005.
- [6] M. E. H. Ismail and X. Li, Bounds for extreme zeros of orthogonal polynomials, Proc. Amer. Math. Soc. 115 (1992), 131–140.
- [7] M. E. H. Ismail and D. R. Masson, q -Hermite polynomials, biorthogonal rational functions, Trans. Amer. Math. Soc. 346 (1994), 63–116.
- [8] M. E. H. Ismail and C. Zhang, Zeros of entire functions and a problem of Ramanujan, Advances in Math., (2007), to appear.
- [9] M. E. H. Ismail and R. Zhang, Scaled asymptotics for q -polynomials, Comptes Rendus, submitted.
- [10] M. E. H. Ismail and R. Zhang, Chaotic and Periodic Asymptotics for q -Orthogonal Polynomials, joint with Mourad E.H. Ismail, International Mathematics Research Notices, accepted.
- [11] R. Koekoek and R. Swarttouw, The Askey-scheme of hypergeometric orthogonal polynomials and its q -analogues, Reports of the Faculty of Technical Mathematics and Informatics no. 98-17, Delft University of Technology, Delft, 1998.
- [12] G. Szegő, Orthogonal Polynomials, Fourth Edition, Amer. Math. Soc., Providence, 1975.
- [13] E. T. Whittaker and G. N. Watson, A Course of Modern Analysis, fourth edition, Cambridge University Press, Cambridge, 1927.
- [14] Ruiming Zhang, Remarks on Ramanujan Function $A_q(z)$, submitted to the Journal of Constructive Approximation.

Current address: School of Mathematics, Guangxi Normal University, Guilin City, Guangxi 541004, P. R. China.

E-mail address: ruimingzhang@yahoo.com